

Haven Hill Acres

Preliminary Drainage Report

JRG Holdings, LLC

518 Leonard Avenue St. Louis, MO 63119 Jimmy

CONSULTING ENGINEER

THOUVENOT, WADE & MOERCHEN, INC. ILLINOIS PROFESSIONAL DESIGN FIRM LICENSE NO. 184-001220 SWANSEA OFFICE 4940 OLD COLLINSVILLE RD. SWANSEA, IL 62226 (618) 624-4488 mmaller@twm-inc.com

Contents

I. Property Overview	1
II. Evaluation of Existing Conditions	
III. Evaluation of Proposed Conditions	
IV. References	

List of Tables

Table 1. Drainage basin characteristics for the existing conditions1

List of Figures

Figure 1. Aerial location map1

Appendices

- A Drainage Area Map & Site Plan B USGS Soils Report

I. Property Overview

The Haven Hill Acres Development is located on parcels 13-1-21-29-12-201-006.002 and 13-1-12-19-12-201-006 in Collinsville, Illinois. The property is currently Zoned R3 Residential in the city of Collinsville. The site is bordered to the east and west by subdivisions zoned R-3. Summit Ridge and Sydney Creek subdivisions border the east. The northern property line is bordered by Reese Drive. The southwest property line is bordered by Carroll Wood Hill Condominiums and the northwest property line is bordered by Sean's Park subdivision. Greenview Acres subdivision is zoned R-3 and is located to the south of the site. A majority of the site is wooded.



Figure 1. Google Earth Aerial location map.

II. Evaluation of Existing Conditions

The site consists of two main drainage areas. Drainage Area #1 drains directly to Schoolhouse Branch to the east. Drainage Area #3 drains to the west through a channel toward the Carroll Wood Hill Subdivision.

FIRM Map 1704390003B, dated 02/18/1981, indicates that the site is listed as Zone C. The soil on the site consists of Menfro, Sylvan, Wakeland and Winfield silt loam. These areas are classified as hydrologic soil group B, indicating moderate infiltration.

Table 1. Drainage basin characteristics for the existing conditions.

Drainage Basin	Area (ac.)	Weighted C
1	15.69	0.20
2	2.99	0.20

III. Evaluation of Proposed Conditions

The proposed construction will include a 10' wide bike path and a multi-family development on a portion of the site. A road connection from Ramada Blvd to Reese Drive is proposed with the development as well. The proposed drainage areas will ultimately drain to various tributaries of Schoolhouse Branch.

Per the City of Collinsville Code of Ordinances, the rational method with rainfall values from Bulletin 75, adjusted for the St. Louis Urban Effect, will be used. The storm sewer network will be sized to pass the 25-year storm event and crossroad culverts will be sized to pass the 50-year storm event. The 5-, 10-, 25-, 50-, and 100-year storm events will be analyzed for the proposed detention pond.

IV. References

City of Collinsville, IL. Code of Ordinances.

Momcilo Markus and J.R. Angel. "Precipitation Frequency Study for Illinois (Bulletin 75)." Illinois State Water Survey.


Illinois Department of Transportation. "Illinois Department of Transportation Drainage Manual." July 2011.

Appendix A

Drainage Area Map & Site Plan



2023\231587\4 CADD - DWG\4.1 Comm\231587-DRAINAGE.dwg Plotted By: bfora



23\231587\4 CADD - DWG\4.1 Comm\231587-DRAINAGE.dwg Plotted By: bf

Appendix B USGS Soils Report



United States Department of Agriculture

Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Madison County, Illinois



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface How Soil Surveys Are Made	
Soil Map	
Soil Map	9
Legend	.10
Map Unit Legend	11
Map Unit Descriptions	.11
Madison County, Illinois	13
79B—Menfro silt loam, 2 to 5 percent slopes	.13
962D2—Sylvan-Bold silt loams, 10 to 18 percent slopes, eroded	14
962F2—Sylvan-Bold silt loams, 18 to 35 percent slopes, eroded	.15
2477B—Winfield-Orthents-Urban land complex, 2 to 8 percent slopes	17
3333A—Wakeland silt loam, 0 to 2 percent slopes, frequently flooded	18
References	.20

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



	MAP LEGEND)	MAP INFORMATION	
Area of In	terest (AOI) Area of Interest (AOI)	8	Spoil Area Stony Spot	The soil surveys that comprise your AOI were mapped at 1:12,000.	
Soils	Soil Map Unit Polygons	03	Very Stony Spot	Warning: Soil Map may not be valid at this scale.	
~	Soil Map Unit Lines Soil Map Unit Points	\$ ∆	Wet Spot Other	Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil	
_	Special Point Features		Special Line Features	line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.	
×	Borrow Pit	Transport	Streams and Canals	Please rely on the bar scale on each map sheet for map	
¥ ♦	Clay Spot Closed Depression	···· ~	Rails Interstate Highways	measurements.	
*	Gravel Pit Gravelly Spot	~	US Routes Major Roads	Source of Map: Natural Resources Conservation Service Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)	
0 1	Landfill Lava Flow	~	Local Roads	Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts	
علله	Marsh or swamp	Backgrou	Background Aerial Photography	distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.	
* 0	Mine or Quarry Miscellaneous Water			This product is generated from the USDA-NRCS certified data as	
0	Perennial Water Rock Outcrop			of the version date(s) listed below. Soil Survey Area: Madison County, Illinois	
+	Saline Spot Sandy Spot			Survey Area Data: Version 19, Aug 21, 2024	
:: =	Severely Eroded Spot			Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.	
♦	Sinkhole Slide or Slip			Date(s) aerial images were photographed: Jul 22, 2022—Aug 25, 2022	
ø	Sodic Spot			The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.	

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
79B	Menfro silt loam, 2 to 5 percent slopes	2.5	13.5%
962D2	Sylvan-Bold silt loams, 10 to 18 percent slopes, eroded	0.2	1.1%
962F2	Sylvan-Bold silt loams, 18 to 35 percent slopes, eroded	13.8	73.8%
2477B	Winfield-Orthents-Urban land complex, 2 to 8 percent slopes	0.2	0.9%
3333A	Wakeland silt loam, 0 to 2 percent slopes, frequently flooded	2.0	10.7%
Totals for Area of Interest		18.7	100.0%

Map Unit Legend

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it

was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Madison County, Illinois

79B—Menfro silt loam, 2 to 5 percent slopes

Map Unit Setting

National map unit symbol: 2x41p Elevation: 390 to 820 feet Mean annual precipitation: 37 to 47 inches Mean annual air temperature: 52 to 57 degrees F Frost-free period: 180 to 200 days Farmland classification: All areas are prime farmland

Map Unit Composition

Menfro and similar soils: 100 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Menfro

Setting

Landform: Hillslopes Landform position (two-dimensional): Summit Landform position (three-dimensional): Side slope Down-slope shape: Convex Across-slope shape: Convex Parent material: Loess

Typical profile

Ap - 0 to 10 inches: silt loam Bt1 - 10 to 33 inches: silty clay loam Bt2 - 33 to 79 inches: silty clay loam

Properties and qualities

Slope: 2 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water supply, 0 to 60 inches: High (about 11.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 2e Hydrologic Soil Group: C Ecological site: F115XB001MO - Deep Loess Upland Woodland Hydric soil rating: No

962D2—Sylvan-Bold silt loams, 10 to 18 percent slopes, eroded

Map Unit Setting

National map unit symbol: 6nzv Elevation: 350 to 1,700 feet Mean annual precipitation: 32 to 41 inches Mean annual air temperature: 48 to 55 degrees F Frost-free period: 150 to 190 days Farmland classification: Not prime farmland

Map Unit Composition

Sylvan and similar soils: 50 percent Bold and similar soils: 40 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sylvan

Setting

Landform: Loess hills on ground moraines Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope Down-slope shape: Linear Across-slope shape: Convex Parent material: Loess

Typical profile

H1 - 0 to 4 inches: silt loam H2 - 4 to 30 inches: silty clay loam H3 - 30 to 80 inches: silt loam

Properties and qualities

Slope: 10 to 18 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 25 percent
Available water supply, 0 to 60 inches: Very high (about 12.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: B Ecological site: F115XC005IL - Loess Upland Forest Hydric soil rating: No

Description of Bold

Setting

Landform: Loess hills on ground moraines Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope Down-slope shape: Linear Across-slope shape: Convex Parent material: Calcareous loess

Typical profile

H1 - 0 to 12 inches: silt loam *H2 - 12 to 60 inches:* silt loam

Properties and qualities

Slope: 10 to 18 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 50 percent
Available water supply, 0 to 60 inches: Very high (about 13.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3e Hydrologic Soil Group: B Ecological site: R115XC002IL - Loess Upland Prairie Hydric soil rating: No

962F2—Sylvan-Bold silt loams, 18 to 35 percent slopes, eroded

Map Unit Setting

National map unit symbol: 6nzw Elevation: 350 to 1,700 feet Mean annual precipitation: 25 to 45 inches Mean annual air temperature: 52 to 57 degrees F Frost-free period: 170 to 200 days Farmland classification: Not prime farmland

Map Unit Composition

Sylvan and similar soils: 50 percent *Bold and similar soils:* 40 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Sylvan

Setting

Landform: Loess bluffs Landform position (two-dimensional): Backslope Down-slope shape: Linear Across-slope shape: Linear

Typical profile

H1 - 0 to 5 inches: silt loam

- H2 5 to 30 inches: silty clay loam
- H3 30 to 80 inches: silt loam

Properties and qualities

Slope: 18 to 35 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 25 percent
Available water supply, 0 to 60 inches: Very high (about 12.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6e Hydrologic Soil Group: B Ecological site: F115XB001MO - Deep Loess Upland Woodland Hydric soil rating: No

Description of Bold

Setting

Landform: Loess bluffs Landform position (two-dimensional): Backslope Down-slope shape: Linear Across-slope shape: Linear

Typical profile

H1 - 0 to 12 inches: silt loam *H2 - 12 to 60 inches:* silt loam

Properties and qualities

Slope: 18 to 35 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 50 percent
Available water supply, 0 to 60 inches: Very high (about 13.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 6e Hydrologic Soil Group: B Ecological site: F114XB804IN - Silty Eolian Forest Hydric soil rating: No

2477B—Winfield-Orthents-Urban land complex, 2 to 8 percent slopes

Map Unit Setting

National map unit symbol: 6qzg Elevation: 330 to 800 feet Mean annual precipitation: 32 to 46 inches Mean annual air temperature: 52 to 57 degrees F Frost-free period: 170 to 200 days Farmland classification: Not prime farmland

Map Unit Composition

Winfield and similar soils: 40 percent Orthents, loamy, and similar soils: 30 percent Urban land: 20 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Winfield

Setting

Landform: Loess hills Landform position (two-dimensional): Summit, shoulder Down-slope shape: Convex Across-slope shape: Convex Parent material: Loess

Typical profile

H1 - 0 to 9 inches: silt loam
H2 - 9 to 13 inches: silt loam
H3 - 13 to 62 inches: silty clay loam
H4 - 62 to 80 inches: silt loam

Properties and qualities

Slope: 2 to 8 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: About 24 to 42 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: High (about 11.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 2e Hydrologic Soil Group: C Ecological site: F115XB001MO - Deep Loess Upland Woodland Hydric soil rating: No

Description of Orthents, Loamy

Setting

Down-slope shape: Convex *Across-slope shape:* Convex

Typical profile

H1 - 0 to 6 inches: loam *H2 - 6 to 60 inches:* loam

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)
Depth to water table: About 42 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: High (about 10.9 inches)

Description of Urban Land

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 8 Hydric soil rating: No

3333A—Wakeland silt loam, 0 to 2 percent slopes, frequently flooded

Map Unit Setting

National map unit symbol: 2wyhj Elevation: 340 to 490 feet Mean annual precipitation: 38 to 49 inches Mean annual air temperature: 50 to 59 degrees F Frost-free period: 180 to 200 days Farmland classification: Prime farmland if drained and either protected from flooding or not frequently flooded during the growing season

Map Unit Composition

Wakeland, frequently flooded, and similar soils: 95 percent

Minor components: 5 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Wakeland, Frequently Flooded

Setting

Landform: Flood plains Landform position (three-dimensional): Talf Down-slope shape: Linear Across-slope shape: Linear Parent material: Silty alluvium

Typical profile

Ap - 0 to 8 inches: silt loam Cg1 - 8 to 19 inches: silt loam Cg2 - 19 to 63 inches: silt loam Cg3 - 63 to 73 inches: silt loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: About 6 to 24 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very high (about 13.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 3w Hydrologic Soil Group: B/D Ecological site: F114XB203IN - Wet Floodplain Forest Hydric soil rating: No

Minor Components

Birds, frequently flooded

Percent of map unit: 5 percent Landform: Flood plains Landform position (three-dimensional): Dip Down-slope shape: Linear Across-slope shape: Linear Ecological site: F113XY919IL - Wet Silty Floodplain Forest Hydric soil rating: Yes

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf